

## Physics 501: Classical Mechanics

Read: **LL** 1-5, **G** 1,2

Special attention: **LL** problems after ch. I

### Homework 1

#### Exercise 1 (G 2.1)

Using variational calculus prove that the shortest distance between two points in space is a straight line.

#### Exercise 2 (G 1.15)

Let  $q_1, \dots, q_n$  be a set of independent generalized coordinates for a system of  $n$  degrees of freedom, with a Lagrangian  $L(q, \dot{q}, t)$ . Suppose we transform to another set of independent coordinates  $s_1, \dots, s_n$  by means of transformation equations

$$q_i = q_i(s_1, \dots, s_n, t), \quad i = 1, \dots, n.$$

(Such a transformation is called a *point transformation*.) Show that if the Lagrangian function is expressed as a function of  $s_j$ ,  $\dot{s}_j$ , and  $t$  through the equations of transformation, then  $L$  satisfies Lagrange's equations with respect to the  $s$  coordinates:

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{s}_j} \right) - \left( \frac{\partial L}{\partial s_j} \right) = 0.$$

In other words, the form of Lagrange's equations is invariant under a point transformation.

*Remark:* Do not use variational principle.

#### Exercise 3

For each of the systems listed below:

- introduce convenient generalized coordinates  $q_j$ ,
- write down the Lagrangian  $L$  as a function of  $q_j$ ,  $\dot{q}_j$  and (if appropriate) time,
- write down the Lagrangian equations of motion.

- a) A particle of the mass  $m$  sliding without friction on a heavy wedge of angle  $\alpha$  and mass  $M$  that can move without friction on a smooth horizontal surface.

- b) A simple pendulum of the mass  $m$  whose point of support moves vertically according to the prescribed law  $y = y_0(t)$ . Pendulum motion is constrained to a vertical plane.
- c) A particle of the mass  $m$  confined to the surface of the sphere (but freely moving along the surface) of radius  $R$  placed in a uniform gravitational field (acceleration  $g$ ).
- d) A bead of the mass  $m$  moving without friction along the rigid wire of elliptic shape  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ .

#### **Exercise 4**

Using variational calculus with Lagrange multiplier, find the shape of the rope of the length  $l$  suspended between two equally high points. The horizontal distance between points is  $2a$ .