

## Homework 4

### Problem 1

A Hamiltonian of some quantum system is given by

$$H = -e^{i\Phi/3}|2\rangle\langle 1| - e^{i\Phi/3}|3\rangle\langle 2| - e^{i\Phi/3}|1\rangle\langle 3| + h.c.,$$

where the states  $|1\rangle$ ,  $|2\rangle$  and  $|3\rangle$  form an orthonormal basis in the Hilbert space of the system,  $\Phi$  is some real number and “*h.c.*” means Hermitian conjugate.

- a) Find the spectrum of this Hamiltonian.
- b) Find the eigenstates of this Hamiltonian.
- c) Find a time-dependent solution  $|\Psi(t)\rangle$  of the Schrodinger equation with this Hamiltonian corresponding to an initial state  $|\Psi(0)\rangle = |1\rangle$ .

### Problem 2

- a) Find the potential profile  $U(x)$  for which the following wavefunction,

$$\Psi(x, t) = Ce^{-ax^2 - ibt}.$$

(with real and constant  $a > 0$  and  $b$ ), satisfies the Schrödinger equation for a particle with mass  $m$ .

- b) Normalize the wavefunction to  $P = 1$ , find  $\langle x \rangle$ ,  $\langle p_x \rangle$ ,  $\delta x$ , and  $\delta p_x$ , and compare the product  $\delta x \delta p_x$  with Heisenberg’s uncertainty relation.

### Problem 3

A quantum particle of the mass  $m$  moves in an infinite potential well  $U(x) = 0$  if  $-a < x < a$  and  $U(x) = \infty$  if  $|x| > a$ .

- a) Write down the spectrum  $E_n$  and normalized eigenfunctions  $\psi_n(x)$ ,  $n = 0, 1, 2, \dots$  for this system.
- b) What is the force exerted by a particle on the walls of the potential well?
- c) Suppose that the particle is initially in the (unnormalized) state  $\psi(x, t = 0) = x$ . What is the probability to find the particle in the ground state ( $n = 0$ ) in time  $t$ ? What is the probability to find the particle in the first excited state  $n = 1$  in time  $t$ ? What is the probability to find this particle in the state  $n = 24$  at the same time?

### Problem 4

A quantum particle of the mass  $m$  moves in a potential well  $U(x) = -U_0$  if  $-a < x < a$  and  $U(x) = 0$  if  $|x| > a$ . Here  $U_0 > 0$  is some constant.

- a) Write down the equations determining the spectrum and eigenstates of bound states in this problem. (the equations are transcendental and can be solved only numerically in this case).
- b) In the limit of a very shallow well  $U_0 \ll \frac{\hbar^2}{ma^2}$  find an approximate energy of the ground state. How many bound states do exist in this case?
- c) Write down an approximate wave function of the ground state in the case of a very shallow well.