

## Homework 15

Problems with stars are not for credit and will NOT be graded.

### Reading

**JJS** 4.3-4.4.

### Problem 1

State the selection rules for the matrix element

$$\langle \alpha'; j', m' | z | \alpha; j, m \rangle,$$

where  $z$  is the component of position operator,  $j, m$  are eigenvalues of  $J^2$  and  $J_z$  respectively and  $\alpha$  are all other quantum numbers.

### Problem 2

The tight binding Hamiltonian on the 1d lattice with periodic boundary conditions is given by

$$H = -W \sum_{n=1}^N \left( e^{i\theta} |n\rangle \langle n+1| + e^{-i\theta} |n+1\rangle \langle n| \right),$$

where  $W$  and  $\theta$  are some real parameters and  $|n\rangle$  form an orthonormal basis of the Hilbert space “state on site  $n$ ” and there is an identification  $|n\rangle \equiv |n+N\rangle$ . This Hamiltonian is obviously translationally invariant (one can change  $n \rightarrow n+1$ ). Use this translational invariance to find the spectrum of the system. Namely,

- a) Write down an operator that translates the system by one unit to the right and check explicitly that it commutes with the Hamiltonian.
- b) Find the spectrum of the translation operator (use  $T^N = 1$ ).
- c) Use previous results to find the spectrum of the Hamiltonian.

### Problem 3

Consider the tight-binding model on three sites

$$H = -W \sum_{n=1}^3 \left( e^{i\theta} |n\rangle \langle n+1| + e^{-i\theta} |n+1\rangle \langle n| \right).$$

Assume that the density matrix of a system at  $t = 0$  is given by

$$\rho(t=0) = \frac{1}{3} (2|1\rangle \otimes \langle 1| + |2\rangle \otimes \langle 2|).$$

Find the density matrix as a function of time  $\rho(t)$ .

#### Problem 4

Assume that some Hamiltonian is symmetric with respect to some discrete group of transformations. The group is generated by two elements  $S$  and  $T$  such that there are relations  $S^2 = 1$ ,  $T^3 = 1$  and  $TST = S$ . For example,  $S$ ,  $T$ ,  $T^2$ ,  $ST$  are all different elements of the group.

- a) What is the spectrum of operator  $S$ ? of operator  $T$ ?
- b) Show that, generally, one expects to find doubly degenerate states in the spectrum of Hamiltonian of the system. What are eigenvalues of  $T$  for those degenerate states? (Hint: consider  $S|\alpha\rangle$  with  $|\alpha\rangle$  being an eigenstate of  $H$ .)
- c) Can you give an example of the system having this symmetry? In your example, what is the meaning of  $S$  and  $T$  symmetry operations?

#### Problem 5

Let  $\phi(\mathbf{p}')$  be the momentum-space wave function for state  $|\alpha\rangle$ , that is,  $\phi(\mathbf{p}') = \langle \mathbf{p}' | \alpha \rangle$ . Is the momentum-space wave function for the time reversed state  $\Theta|\alpha\rangle$  given by  $\phi(\mathbf{p}')$ ,  $\phi(-\mathbf{p}')$ ,  $\phi^*(\mathbf{p}')$ , or  $\phi^*(-\mathbf{p}')$ ? Justify your answer.

#### Problem 6

The Hamiltonian for a spin 1 system is given by

$$H = AS_z^2 + B(S_x^2 - S_y^2).$$

Solve this problem *exactly* to find the normalized energy eigenstates and eigenvalues. (A spin-dependent Hamiltonian of this kind actually appears in crystal physics.) Is this Hamiltonian invariant under time reversal? How do the normalized eigenstates you obtained transform under time reversal?